

The spectral theorem of many-body Green's function theory when there are zero eigenvalues of the matrix governing the equations of motion.

P. Fröbrich* and P.J. Kuntz†

Hahn-Meitner-Institut Berlin, Glienicker Straße 100, D-14109 Berlin, Germany

(Dated: February 2, 2008)

In using the spectral theorem of many-body Green's function theory in order to relate correlations to commutator Green's functions, it is necessary in the standard procedure to consider the anti-commutator Green's functions as well whenever the matrix governing the equations of motion for the commutator Green's functions has zero eigenvalues. We show that a singular-value decomposition of this matrix allows one to reformulate the problem in terms of a smaller set of Green's functions with an associated matrix having no zero eigenvalues, thus eliminating the need for the anti-commutator functions. The procedure is quite general and easy to apply. It is illustrated for the field-induced reorientation of the magnetization of a ferromagnetic Heisenberg monolayer and it is expected to work for more complicated cases as well.

PACS numbers: 75.10.Jm, 75.70.Ak

Many-body Green's function (GF) techniques make use of the spectral theorem to connect correlation functions to the Green's functions, which in turn are determined from equations of motion that can be formulated conveniently in terms of a matrix equation. Should this matrix have zero eigenvalues, particular care must be taken, since they cannot be handled directly by the spectral theorem for the commutator GF, since it would require a division by zero. The standard treatment of this case demands in addition to the commutator GF a knowledge of the anti-commutator GF (see references [1, 2] and textbooks [3, 4]). An equivalent treatment in terms of the anti-commutator GF alone may formally appear to avoid this problem, but in practice one is faced with the same numerical difficulties. In this paper, we show that these difficulties can be overcome by using a singular-value decomposition (SVD) of the equations of motion matrix to eliminate the space connected with the zero eigenvalues (null-space), thus reducing the size of the problem and, at the same time, making the anti-commutator GF superfluous. That is, one uses projection matrices furnished by the SVD to reformulate the spectral theorem in terms of a smaller number of commutator GF's whose associated matrix has no zero eigenvalues. We show that this procedure is generally applicable and offers an alternative to the standard treatment; i.e. we show that a knowledge of the commutator GF's is sufficient.

The appearance of zero eigenvalues in the matrix governing the equations of motion is *not* an unusual event. We have met this problem in much of our previous work on the application of many-body GF's to the description of thin ferromagnetic films [5, 6, 7, 8, 9, 10]. We treated the Heisenberg ferromagnet including single-ion anisotropy and exchange anisotropy for several values of the spin S for single-layer and multi-layer systems. A variety of decoupling approximations [11, 12] to arrive at a closed set of equations of motion were considered. In most of these cases, the equations of motion matrix

has at least one zero eigenvalue per layer and, if higher-order Green's functions [7] are used to achieve an exact treatment of the single-ion anisotropy, there are two zero eigenvalues per layer. Calculation of many-layer systems therefore leads to matrices with a large number of zero eigenvalues. This, coupled with the fact that the matrices are in general nonsymmetric, leads in some cases to numerical difficulties which can best be circumvented by eliminating the null-space. The method for doing this may be considered as an alternative to the standard procedure of handling zero eigenvalues. In the following, we first introduce our methodology and notation by reviewing briefly the standard text-book treatment of zero eigenvalues, then we describe the new method which is based on the singular value decomposition of the matrix governing the equation of motion, and finally, by way of example, we apply the new method to the field-induced reorientation of the magnetization of a ferromagnetic Heisenberg monolayer.

The standard procedure.—The equation of motion for a Green's function vector

$$\mathbf{G}_{\eta,\omega} = \langle\langle A; B \rangle\rangle_{\eta,\omega} \quad (1)$$

in energy space is (omitting the subscript ω)

$$\omega \langle\langle A; B \rangle\rangle_{\eta} = \langle[A, B]_{\eta}\rangle + \langle\langle[A, H]_{-}; B\rangle\rangle_{\eta}, \quad (2)$$

where $\langle\ldots\rangle = \text{Tr}(\ldots e^{-\beta H})/Z$ is the thermodynamic expectation value, H is the Hamiltonian and A and B are operators depending on the problem under consideration, and $\eta = \mp 1$ refers to the commutator or anti-commutator Green's functions, respectively.

Repeated application of equation (2) to the higher-order Green's functions appearing on the right hand side, $\langle\langle[A, H]_{-}; B\rangle\rangle$, results in an infinite hierarchy of equations. In order to obtain a set of solvable equations, one has to terminate this hierarchy, usually by expressing the Green's functions of a particular order in terms of those of

lower order. In many applications, only the lowest-order functions are retained using a decoupling procedure of the form

$$\langle\langle[A, H]_-; B\rangle\rangle_\eta \simeq \mathbf{\Gamma}\langle\langle A; B\rangle\rangle_\eta, \quad (3)$$

where $\mathbf{\Gamma}$ is the (in general *non-symmetric*) matrix expressing the higher-order Green's functions in terms of linear combinations of lower-order ones. In compact matrix notation, the equation of motion is

$$(\omega \mathbf{1} - \mathbf{\Gamma})\mathbf{G}_\eta = \mathbf{A}_\eta, \quad (4)$$

where $\mathbf{1}$ is the unit matrix, and the inhomogeneity vector is $\mathbf{A}_\eta = \langle[A, B]_\eta\rangle$.

It is now convenient to introduce the notation of the eigenvector method of reference [9], since it is particularly suitable for the multi-dimensional problems in which many zero eigenvalues are likely to appear. One starts by diagonalizing the matrix $\mathbf{\Gamma}$

$$\mathbf{L}\mathbf{R} = \mathbf{\Omega}, \quad (5)$$

where $\mathbf{\Omega}$ is the diagonal matrix of N eigenvalues, ω_τ ($\tau = 1, \dots, N$), N_0 of which are zero and $(N - N_0)$ are non-zero. The matrix $\mathbf{R}$ contains the right eigenvectors as columns and its inverse $\mathbf{L} = \mathbf{R}^{-1}$ contains the left eigenvectors as rows. $\mathbf{L}$ is constructed such that $\mathbf{L}\mathbf{R} = \mathbf{R}\mathbf{L} = \mathbf{1}$. Multiplying equation (4) from the left by $\mathbf{L}$, inserting $\mathbf{1} = \mathbf{R}\mathbf{L}$, and defining new vectors $\mathcal{G}_\eta = \mathbf{L}\mathbf{G}_\eta$ and $\mathcal{A}_\eta = \mathbf{L}\mathbf{A}_\eta$ one finds

$$(\omega \mathbf{1} - \mathbf{\Omega})\mathcal{G}_\eta = \mathcal{A}_\eta. \quad (6)$$

Each of the components τ of this Green's function vector has but a single pole

$$(\mathcal{G}_\eta)_\tau = \frac{(\mathcal{A}_\eta)_\tau}{\omega - \omega_\tau}. \quad (7)$$

This allows the spectral theorem, (see e.g. [3, 4]), to be applied to each component of the Greens's function vector *separately*. Define the correlation vector $\mathcal{C} = \mathbf{L}\mathbf{C}$, where $\mathbf{C} = \langle BA \rangle$ is the vector of correlations associated with $\mathbf{G}_\eta$. From the spectral theorem, the component τ for the anti-commutator case ($\eta = +1$) is

$$C_\tau = \frac{(\mathcal{A}_\eta)_\tau}{e^{\beta\omega_\tau} + \eta}. \quad (8)$$

Use of the anti-commutator functions appears at first sight more straightforward (zero eigenvalues seem to make no problem), but leads in general to momentum dependent inhomogeneities, which are difficult to treat. However, it is usually easier to employ commutator Green's functions, because in this case the inhomogeneity vectors are momentum independent.

For the commutator functions ($\eta = -1$), the $N - N_0$ components for $\omega_\tau \neq 0$, $\mathcal{C}^1$ (the upper index 1 refers to the non-null space), are

$$(\mathcal{C}^1)_\tau = \frac{(\mathcal{A}_\eta)_\tau}{e^{\beta\omega_\tau} + \eta}. \quad (9)$$

This equation cannot be used to define the N_0 components $\mathcal{C}^0$ corresponding to $\omega_\tau = 0$ because of the zero in the denominator. Instead, one must enlist the help of the *anti-commutator* Green's function:

$$(\mathcal{C}^0)_{\tau_0} = \lim_{\omega \rightarrow 0} \frac{1}{2} \omega (\mathcal{G}_{\eta=+1})_\tau. \quad (10)$$

The components of $\mathcal{C}^0$, indexed by τ_0 (the upper index 0 refers to the null space), can be simplified by using the relation between the commutator and anti-commutator inhomogeneities, $\mathbf{A}_1 = \mathbf{A}_{-1} + 2\mathbf{C}$. This yields, for $\omega_{\tau_0} = 0$,

$$\begin{aligned} (\mathcal{C}^0)_{\tau_0} &= \frac{1}{2} \lim_{\omega \rightarrow 0} \frac{\omega (\mathcal{A}_{+1})_{\tau_0}}{\omega - \omega_{\tau_0}} = \frac{1}{2} (\mathcal{A}_{+1})_{\tau_0} \\ &= \frac{1}{2} (\mathbf{L}^0 (\mathbf{A}_{-1} + 2\mathbf{C}))_{\tau_0} = (\mathbf{L}^0 \mathbf{C})_{\tau_0}, \end{aligned} \quad (11)$$

where use has been made of the relation $\mathbf{L}^0 \mathbf{A}_{-1} = 0$, which derives from the fact that the commutator Green's function is regular at the origin [9].

At this point it is again convenient to introduce superscripts 0 and 1 to denote vectors corresponding to zero and non-zero eigenvalues, respectively. The right and left eigenvectors and the correlation vectors may then be partitioned as $\mathbf{R} = (\mathbf{R}^1 \mathbf{R}^0)$ and

$$\mathbf{L} = \begin{pmatrix} \mathbf{L}^1 \\ \mathbf{L}^0 \end{pmatrix} \quad \mathcal{C} = \begin{pmatrix} \mathcal{C}^1 \\ \mathcal{C}^0 \end{pmatrix}, \quad (12)$$

where the correlation vectors from equations (9) and (11) are then $\mathcal{C}^0 = \mathbf{L}^0 \mathbf{C}$ and $\mathcal{C}^1 = \mathcal{E}^1 \mathbf{L}^1 \mathbf{A}$, and $\mathcal{E}^1$ is the $(N - N_0) \times (N - N_0)$ matrix with $1/(e^{\beta\omega_\tau} - 1)$ on the diagonal.

Multiplying the correlation vector $\mathcal{C}$ from the left by $\mathbf{R}$ yields a compact matrix equation for the original correlation vector $\mathbf{C}$:

$$\mathbf{C} = \mathbf{R}^1 \mathcal{E}^1 \mathbf{L}^1 \mathbf{A} + \mathbf{R}^0 \mathbf{L}^0 \mathbf{C}, \quad (13)$$

This equation is in momentum space and the coupled system of integral equations obtained by Fourier transformation to real space has to be solved self-consistently. This was done in all our previous work [5, 6, 7, 8, 9, 10].

The singular value decomposition.— We now present an alternative method of treating the zero eigenvalues. Starting again from the equation of motion (4), we write the $\mathbf{\Gamma}$ matrix as a singular value decomposition [13]:

$$\mathbf{\Gamma} = \mathbf{U}\mathbf{W}\tilde{\mathbf{V}}. \quad (14)$$

The matrix $\mathbf{W}$ is a diagonal matrix whose elements are the singular values, which are ≥ 0 and $\mathbf{U}$ and $\mathbf{V}$ are orthogonal matrices: $\tilde{\mathbf{U}}\mathbf{U} = \mathbf{1}$ and $\tilde{\mathbf{V}}\mathbf{V} = \mathbf{1}$, where $\tilde{\mathbf{V}}$ denotes the transpose of $\mathbf{V}$. $\mathbf{U}$, $\mathbf{V}$ and $\mathbf{W}$ can be obtained very efficiently numerically [13]. However, one can also get the vectors $\mathbf{V}$ and $\mathbf{U}$ directly by diagonalizing $\tilde{\mathbf{\Gamma}}\mathbf{\Gamma}$ or $\mathbf{\Gamma}\tilde{\mathbf{\Gamma}}$ respectively. The singular values are just the positive square roots of the eigenvalues of these matrices, $\mathbf{W}^2$.

$$\begin{aligned}\tilde{\mathbf{V}}\tilde{\mathbf{\Gamma}}\mathbf{\Gamma}\mathbf{V} &= \tilde{\mathbf{V}}\mathbf{V}\mathbf{W}\tilde{\mathbf{U}}\mathbf{U}\mathbf{W}\tilde{\mathbf{V}} = \mathbf{W}^2, \\ \tilde{\mathbf{U}}\tilde{\mathbf{\Gamma}}\mathbf{\Gamma}\mathbf{U} &= \tilde{\mathbf{U}}\mathbf{U}\mathbf{W}\tilde{\mathbf{V}}\mathbf{V}\mathbf{W}\tilde{\mathbf{U}} = \mathbf{W}^2.\end{aligned}\quad (15)$$

Note that the $\mathbf{\Gamma}$ matrix is fully determined by the non-zero singular values and the corresponding eigenvectors:

$$\mathbf{\Gamma} = \mathbf{U}\mathbf{W}\tilde{\mathbf{V}} = \mathbf{u}\mathbf{w}\tilde{\mathbf{v}}. \quad (16)$$

$\mathbf{u}$ and $\mathbf{v}$ are $N \times (N - N_0)$ matrices obtained from $\mathbf{U}$ and $\mathbf{V}$ by omitting the columns corresponding to the zero singular values. The matrix $\mathbf{w}$ is the $(N - N_0) \times (N - N_0)$ diagonal matrix with the non-zero singular values on the diagonal.

The crucial point is that the dimension of the equations of motion can be reduced by the number of zero singular values by applying the following transformations

$$\begin{aligned}\gamma &= \tilde{\mathbf{v}}\mathbf{\Gamma}\mathbf{v}, \\ \mathbf{g} &= \tilde{\mathbf{v}}\mathbf{G}, \\ \mathbf{a} &= \tilde{\mathbf{v}}\mathbf{A},\end{aligned}\quad (17)$$

and realizing that $\mathbf{\Gamma} = \mathbf{\Gamma}v\tilde{v}$ and $\tilde{v}v = 1$.

The $(N - N_0)$ -dimensional equation of motion then becomes

$$(\omega\mathbf{1} - \gamma)\mathbf{g} = \mathbf{a}. \quad (18)$$

Here again the eigenvector method can be used, where $\mathbf{l}$ and $\mathbf{r}$ diagonalize the γ -matrix, $\mathbf{l}\gamma\mathbf{r} = \omega^1$. Applying the spectral theorem to the equation of motion with the matrix γ (which now has no zero eigenvalues) yields the equation for the correlations $\mathbf{c} = \tilde{\mathbf{v}}\mathbf{C}$ in momentum space

$$\mathbf{c} = \mathbf{r}\mathcal{E}^1\mathbf{la} \quad (19)$$

where the $(N - N_0) \times (N - N_0)$ diagonal matrix $\mathcal{E}^1$ has the same elements as before, $1/(e^{\beta\omega_\tau} - 1)$. In order to determine the correlations in coordinate space, one has to perform a Fourier transform and then solve self-consistently the system of integral equations

$$0 = \int d\mathbf{k}(\mathbf{r}\mathcal{E}^1\mathbf{l}\tilde{\mathbf{v}}\mathbf{A} - \tilde{\mathbf{v}}\mathbf{C}), \quad (20)$$

where $\mathbf{A}$ and $\mathbf{C}$ are the inhomogeneity and correlation vectors of the original problem, and the integration is over the first Brillouin zone.

The advantages of the new procedure are that, after projecting onto the non-null space, it is no longer necessary to calculate the term in the spectral theorem which is connected with the anti-commutator Green's function, and the dimension of the final system of equations is reduced by the number of zero eigenvalues of the original $\mathbf{\Gamma}$ matrix.

Example: The Heisenberg monolayer.— As an example, we apply the new procedure to a ferromagnetic Heisenberg monolayer with spin $S = 1$, in which the magnetization can be reoriented by applying an external field. This is an example in which zero eigenvalues of the $\mathbf{\Gamma}$ matrix occur and is treated in references [6, 9] with the standard procedure not only for the $S = 1$ monolayer but also for the multilayer case and for $S \geq 1$.

We restrict ourselves to a spin Hamiltonian consisting of an isotropic Heisenberg exchange interaction with strength J_{kl} between nearest neighbour lattice sites, a second-order single-ion lattice anisotropy with strength $K_{2,k}$, and an external magnetic field $\mathbf{B} = (B^x, B^y, B^z)$:

$$\begin{aligned}\mathcal{H} = & - \frac{1}{2} \sum_{\langle kl \rangle} J_{kl} (S_k^- S_l^+ + S_k^z S_l^z) - \sum_k K_{2,k} (S_k^z)^2 \\ & - \sum_k \left(\frac{1}{2} B^- S_k^+ + \frac{1}{2} B^+ S_k^- + B^z S_k^z \right).\end{aligned}\quad (21)$$

Here the notation $S_k^\pm = S_k^x \pm iS_k^y$ and $B^\pm = B^x \pm iB^y$ is introduced, where k and l are lattice site indices and $\langle kl \rangle$ indicates summation over nearest neighbours only.

Applying a Tyablikov (RPA) decoupling to the exchange interaction term, and an Anderson-Callen decoupling to the single-ion anisotropy term one obtains [6, 9] equations of motion for the monolayer of the form

$$(\omega\mathbf{1} - \mathbf{\Gamma})\mathbf{G}_\eta = \mathbf{A}_\eta \quad (22)$$

with the Green's function and inhomogeneity vectors

$$\mathbf{G}_\eta^{mn}(\mathbf{k}, \omega) = \begin{pmatrix} G_\eta^{+,mn}(\mathbf{k}, \omega) \\ G_\eta^{-,mn}(\mathbf{k}, \omega) \\ G_\eta^{z,mn}(\mathbf{k}, \omega) \end{pmatrix}, \quad \mathbf{A}_\eta^{mn} = \begin{pmatrix} A_\eta^{+,mn} \\ A_\eta^{-,mn} \\ A_\eta^{z,mn} \end{pmatrix}, \quad (23)$$

where $G_{ij,\eta}^{\alpha,mn}(\omega) = \langle\langle S_i^\alpha; (S_j^z)^m (S_j^-)^n \rangle\rangle_{\omega,\eta}$ and $\alpha = (+, -, z)$ takes care of all directions in space. For a field in the x -direction only, $\mathbf{B} = (B^x, 0, 0)$, the reorientation of the magnetization takes place in the xz -plane, and the $\mathbf{\Gamma}$ matrix for spin $S = 1$ ($n \geq 1, m \geq 0, m + n \leq 2$) is

$$\mathbf{\Gamma} = \begin{pmatrix} H^z & 0 & -H^x \\ 0 & -H^z & H^x \\ -\frac{1}{2}H^x & \frac{1}{2}H^x & 0 \end{pmatrix}. \quad (24)$$

where

$$\begin{aligned}H^z &= Z + \langle S^z \rangle J(q - \gamma_{\mathbf{k}}), \\ Z &= K_2 2 \langle S^z \rangle \left(1 - \frac{1}{2S^2} [S(S+1) - \langle S^z S^z \rangle] \right), \\ H^x &= B^x + \langle S^x \rangle J(q - \gamma_{\mathbf{k}}).\end{aligned}\quad (25)$$




FIG. 1: Magnetizations $\langle S^z \rangle$ and $\langle S^x \rangle$ and the reorientation angle Θ for a spin $S=1$ Heisenberg monolayer as function of the temperature.

For a square lattice with lattice constant unity, $\gamma_{\mathbf{k}} = 2(\cos k_x + \cos k_y)$, and $q = 4$ is the number of intra-layer nearest neighbours. The singular value decomposition of equation (14) can be performed analytically for the monolayer case. The singular values of the diagonal matrix W are given as the positive square roots of the eigenvalues of the matrix $\mathbf{\Gamma}\tilde{\mathbf{\Gamma}}$ or $\tilde{\mathbf{\Gamma}}\mathbf{\Gamma}$

$$\mathbf{W} = \begin{pmatrix} \epsilon_1 & 0 & 0 \\ 0 & \epsilon_2 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad (26)$$

where $\epsilon_1 = \sqrt{H^z H^z + 2H^x H^x}$, $\epsilon_2 = \sqrt{H^z H^z + \frac{1}{2}H^x H^x}$, and one singular value zero. The matrices $\mathbf{U}$ and $\mathbf{V}$ are obtained from the eigenvectors of these matrices

$$\mathbf{U} = \begin{pmatrix} \frac{-\sqrt{2}}{2} & \frac{-H^z}{\sqrt{2}\epsilon_2} & \frac{H^x}{2\epsilon_2} \\ \frac{\sqrt{2}}{2} & \frac{-H^z}{\sqrt{2}\epsilon_2} & \frac{H^x}{2\epsilon_2} \\ 0 & \frac{\sqrt{2}\epsilon_2}{H^z} & \frac{H^z}{\epsilon_2} \end{pmatrix}. \quad (27)$$

and

$$\tilde{\mathbf{V}} = \begin{pmatrix} \frac{-H^z}{\sqrt{2}\epsilon_1} & \frac{-H^z}{\sqrt{2}\epsilon_1} & \frac{\sqrt{2}H^x}{\epsilon_1} \\ \frac{-\sqrt{2}}{2} & \frac{\sqrt{2}}{2} & 0 \\ \frac{H^x}{\epsilon_1} & \frac{H^x}{\epsilon_1} & \frac{H^z}{\epsilon_1} \end{pmatrix}. \quad (28)$$

The lower case rectangular matrices $\mathbf{u}$ and $\tilde{\mathbf{v}}$ introduced above are obtained by omitting the last column in $\mathbf{U}$ and the last row in $\tilde{\mathbf{V}}$ respectively. One sees explicitly that $\tilde{\mathbf{v}}\mathbf{v} = \mathbf{1}$ and that $\mathbf{v}\tilde{\mathbf{v}}$ is a projection operator onto the non-null space because it has two eigenvalues equal to 1 and one equal to 0, and $(\mathbf{v}\tilde{\mathbf{v}})^2 = \mathbf{v}\tilde{\mathbf{v}}$.

Having constructed the matrix $\tilde{\mathbf{v}}$, we now solve equation (20) for the spin $S = 1$ monolayer with the curve following method, described in detail in reference [7]. The

result, shown in figure 1, agrees perfectly with that obtained in references [6, 9] obtained with the standard procedure, i.e. by solving equation (13). In that work, spins $S > 1$ and multilayer systems were also considered. This can also be handled by the new algorithm but, in this case, the singular value decomposition and the matrix $\tilde{\mathbf{v}}$ have to be obtained from an appropriate numerical method [13].

Summary.— We have developed a new method for dealing with zero eigenvalues of the matrix governing the equations of motion for many-body Green's functions by applying a singular value decomposition to this matrix. This enables one to eliminate the null space, thus reducing the number of integral equations which have to be solved self-consistently by the number of zero eigenvalues of the equation of motion matrix. The calculation of the anti-commutator Green's function, necessary in the standard procedure using the spectral theorem, is superfluous in the new procedure, which is demonstrated for the example of the field induced reorientation of the magnetization of a spin $S = 1$ Heisenberg ferromagnetic monolayer. The method is expected to work for more complicated cases, where it has decided advantages over the usual procedure.

* Also at Institut für Theoretische Physik, Freie Universität Berlin, Arnimallee 14, D-14195 Berlin, Germany.; Electronic address: froebrich@hmi.de

† Electronic address: kuntz@hmi.de

- [1] K.W.H. Stevens, G.A. Tombs, Proc. Phys. Soc. **85**, 1307 (1965).
- [2] J.G. Ramos, A.A. Gomes, Il Nuovo Cimento **3**, 441 (1971).
- [3] W. Nolting, Quantentheorie des Magnetismus, vol.2, Stuttgart: Teubner, 1986, Chapter B.1.3.
- [4] W. Gasser, E. Heiner, and K. Elk, in 'Greensche Funktionen in der Festkörper- und Vielteilchenphysik', Wiley-VHC, Berlin, 2001, Chapter 3.3.
- [5] A. Ecker, P. Fröbrich, P.J. Jensen, P.J. Kuntz, J. Phys.: Condens. Matter **11**, 1557 (1999).
- [6] P. Fröbrich, P.J. Jensen, P.J. Kuntz, Eur. Phys. J. B **13**, 477 (2000).
- [7] P. Fröbrich, P.J. Kuntz, M. Saber, Ann. Phys. (Leipzig) **11**, 387 (2002).
- [8] P. Henelius, P. Fröbrich, P.J. Kuntz, C. Timm, P.J. Jensen, Phys. Rev. B **66**, 094407 (2002).
- [9] P. Fröbrich, P.J. Jensen, P.J. Kuntz, A. Ecker, Eur. Phys. J. B **18**, 579 (2000).
- [10] P. Fröbrich, P.J. Kuntz, in preparation
- [11] S.V. Tyablikov, Ukr. Mat. Zh. **11**, 289 (1959).
- [12] F.B. Anderson, H.B. Callen, Phys. Rev. **136**, A1068 (1964).
- [13] W.H. Press, B.P. Flannery, S.A. Teukolsky, W.T. Vetterling, Numerical Recipes, Cambridge University Press, 1989.